

## 学位論文の要旨

# Lubiprostone Decreases the Small Bowel Transit Time by Capsule Endoscopy: An Exploratory, Randomised, Double-Blind, Placebo-Controlled 3-Way Crossover Study (健康成人男性におけるルビプロストンによる胃・小腸通過時間への影響) (-カプセル内視鏡を用いて-)

Mizue Matsuura

松浦 瑞恵

Hepatology and Gastroenterology  
Yokohama City University Graduate School of Medicine  
横浜市立大学 大学院医学研究科 医科学専攻 肝胆膵消化器病学

( Doctoral Supervisor : Atsushi Nakajima, Professor )

( 指導教員 : 中島 淳 教授 )

# Lubiprostone Decreases the Small Bowel Transit Time by Capsule Endoscopy: An Exploratory, Randomised, Double-Blind, Placebo-Controlled 3-Way Crossover Study

(健康成人男性におけるルビプロストンによる胃・小腸通過時間への影響)  
(-カプセル内視鏡を用いて-)

<http://www.hindawi.com/journals/grp/2014/879595/>

## Introduction:

Lubiprostone (Amitiza; Takeda Pharmaceuticals North America, Deerfield, IL) selectively activates the type-2 chloride channels in the apical membrane of the GI epithelium, inducing net fluid secretion. The proposed primary mechanism of action of lubiprostone in the gastrointestinal tract is increased chloride ion transport into the intestinal lumen by the drug caused by the opening of ClC-2, which results in increased intestinal secretion and accelerated mass transit [J. Cuppoletti et al., 2004. and A. Postgate et al., 2009.]. The aim of this study was to investigate the usefulness of lubiprostone for bowel preparation and as a propulsive agent in small bowel endoscopy.

## Methods:

This was a randomized, double-blind, placebo-controlled, 3-way crossover study of subjects who volunteered to undergo CE. The subjects were 6 asymptomatic male volunteers (average age: 35.8 years; age range: 29–50 years). In all the subjects, the CE was performed with the PillCam SB2 CE system (Given Imaging Ltd.), and the images were viewed with the Rapid 5 Reader. The subjects were randomly assigned to receive a 24  $\mu$ g tablet of lubiprostone 60 minutes prior to the capsule ingestion for CE and a placebo tablet 30 minutes before the capsule ingestion (L-P regimen), a placebo tablet 60 minutes prior to the capsule ingestion for CE and a 24  $\mu$ g tablet of lubiprostone 30 minutes prior to the capsule ingestion (P-L regimen), or a placebo tablet 60 minutes prior to the capsule ingestion for CE and a placebo tablet again 30 minutes prior to the capsule ingestion (P-P regimen). Each of the test conditions was separated by a washout period of at least 7 days.

All the study subjects were instructed to have a light breakfast and then clear liquids on the day prior to the CE. Furthermore, they were instructed to have nothing by mouth for at least 8 hours prior to the capsule ingestion for CE. All the CE images were read by two investigators (Masahiko Inamori and Mizue Matsuura), both of whom were blinded to the group allocation status of the

subjects. The small bowel examination was considered to be complete if the capsule had passed into the colon.

The quality of the capsule endoscopic images and the amount of water in the small bowel were assessed on 5-point scale. We used a 5-point scale (0–4) based on the percentage of the capsule images that were unimpaired by the presence of debris or dark luminal fluid (4, 100–80%; 3, 80–60%; 2, 60–40%; 1, 40–20%; 0, 20–0%). The average scores for 5min segments of the video were assessed from capsule entry into the proximal duodenum (0% of the SBTT) and for every 10% of the SBTT thereafter, with the score for the final segment recorded in the terminal ileum (100% of the SBTT).

### **Results:**

The study was completed in six male subjects (mean age: 39.5 years; range: 29–50 years). No adverse events occurred during the study. The median GTT was 22.5 (9–160) minutes in the P-P regimen, 40 (4–122) minutes in the L-P regimen, and 57.5 (15–78) minutes in the P-L regimen ( $P = 0.846$ ). The median SBTT was 178.5 (117–407) minutes in the P-P regimen, 122.5 (27–282) minutes in the L-P regimen, and 110.5 (11–331) minutes in the P-L regimen ( $P = 0.042$ ). The median SBTT values for the L-P and P-L regimens were statistically significantly different from the SBTT in the P-P regimen.

The image quality score was  $2.88 \pm 1.35$  in the P-P regimen,  $3.56 \pm 0.56$  in the L-P regimen, and  $3.76 \pm 0.85$  in the P-L regimen ( $P < 0.001$ ). The amount of water in the small bowel was  $1.66 \pm 1.65$  in the P-P regimen,  $3.13 \pm 1.64$  in the L-P regimen, and  $2.60 \pm 1.29$  in the P-L regimen ( $P < 0.01$ ).

### **Discussion:**

This study was designed to evaluate the effect of lubiprostone on the capsule transit time through the GI lumen and its effectiveness as a bowel preparation agent for improving the quality of capsule imaging of the small bowel. Lubiprostone improved the imaging quality of the small bowel as compared to placebo and also improved the SBTT. The GTT following administration of lubiprostone was similar to that after administration of placebo. Our findings differ from those of the study reported by Camilleri et al. 2006, who reported finding evidence of delayed gastric emptying following the administration of lubiprostone. The main side effect of lubiprostone was nausea, possibly related to delayed gastric emptying. Lubiprostone improved the SBTT as compared to placebo. Camilleri et al. showed lubiprostone decreased SBTT in 2006. In addition, lubiprostone has been shown to accelerate overall colonic transit without significantly changing the rate of emptying of the ascending colon [J. Cuppoletti et al., 2004.]. Our study shows that lubiprostone significantly decreases the SBTT and improves the visualization of the small bowel during CE. We also confirm that lubiprostone induces water secretion into the small bowel lumen by capsule endoscopy.

## References

Camilleri M, Bharucha AE, Ueno R, Burton D, Thomforde GM, Baxter K, McKinzie S, Zinsmeister AR., (2006) "Effect of a selective chloride channel activator, lubiprostone, on gastrointestinal transit, gastric sensory, and motor functions in healthy volunteers," *American Journal of Physiology:Gastrointestinal and Liver Physiology*, vol. 290, no. 5, pp. G942–G947.

Carey EJ, Leighton JA, Heigh RI, Shiff AD, Sharma VK, Post JK, Fleischer DE., (2006) "A single-center experience of 260 consecutive patients undergoing capsule endoscopy for obscure gastrointestinal bleeding." *Am J Gastroenterol* ,102, 89–95.

Cryer B, (2014) "A randomized study of lubiprostone for opioid-induced constipation in patients with chronic noncancer pain." *Pain Med.*15(11),1825-34.

Cuppoletti J, Malinowska DH, Tewari KP, Li QJ, Sherry AM, Patchen ML, Ueno R., (2004) "SPI-0211 activates T84 cell chloride transport and recombinant human ClC-2 chloride currents," *The American Journal of Physiology—Cell Physiology*, vol. 287, no. 5, pp. C1173–C1183.

Hooks SB 3rd, Rutland TJ, Di Palma JA, (2009) "Lubiprostone neither decreases gastric and small-bowel transit time nor improves visualization of small bowel for capsule endoscopy: a double-blind, placebo-controlled study." *Gastrointest Endoscopy*. Nov;70(5):942-6.

Hosono K, Endo H, Sakai E, Sekino Y, Uchiyama T, Watanabe S, Iida H, Sakamoto Y, Koide T, Takahashi H, Yoneda M, Tokoro C, Abe Y, Inamori M, Kobayashi N, Kubota K, Nakajima A., (2011) "Optimal Approach for Small Bowel Capsule Endoscopy Using Polyethylene Glycol and Metoclopramide with the Assistance of a Real-Time Viewer." *Digestion*, 84(2):119-25.

Johanson JF, Ueno R., (2007) "Lubiprostone, a locally acting chloride channel activator, in adult patients with chronic constipation: a double-blind, placebo-controlled, dose-ranging study to evaluate efficacy and safety." *Alimentary Pharmacology & Therapeutics*, vol. 25.

Postgate A, Tekkis P, Patterson N, Fitzpatrick A, Bassett P, Fraser C., (2009) "Are bowel purgatives and prokinetics useful for small-bowel capsule endoscopy? A prospective randomized controlled study," *Gastrointestinal Endoscopy*, vol. 69, no. 6, pp. 1120–1128.

Rondonotti E, Herrerias JM, Pennazio M, Caunedo A, Mascarenhas-Saraiva M, de Franchis R., (2005) "Complications, limitations, and failures of capsule endoscopy: a review of 733 cases." *Gastrointest Endosc.*, 62

Sharma A, Jamal MM., (2013) "Opioid induced bowel disease: a twenty-first century physicians' dilemma." *Curr Gastroenterol.Rep.*15:334.

Sykes N., (2010) "Constipation and diarrhea. *In: Oxford Textbook of Palliative Medicine*", 4<sup>th</sup> ed. *Oxford University Press, New York*, pp.833-850.

Tatar EL, Shen EH, Palance AL, Sun JH, Pitchumoni CS., (2006) "Clinical utility of wireless capsule endoscopy: experience with 200 cases." *J Clin Gastroenterol* , 40, 140–144.

Thomas J., (2007) "Cancer-related constipation." *Curr. Oncol. Rep.*, 9:278-284.

## 論文目録

### 主論文

“Lubiprostone Decreases the Small Bowel Transit Time by Capsule Endoscopy: An Exploratory, Randomised, Double-Blind, Placebo-Controlled 3-Way Crossover Study”

**MizueMatsuura**, Masahiko Inamori, Hiroki Endo, Tetsuya Matsuura, Kenji Kanoshima, Yumi Inoh, Yuji Fujita, Shotaro Umezawa, Akiko Fuyuki, Shiori Uchiyama, Takuma Higurashi, Hidenori Ohkubo, Eiji Sakai, Hiroshi Iida, Takashi Nonaka, Seiji Futagami, Akihiko Kusakabe, Shin Maeda, and Atsushi Nakajima.

*Gastroenterol Res Pract.* 2014;2014:879595. doi: 10.1155/2014/879595. Epub 2014 Dec 29.

### 副論文

“Effectiveness of therapeutic barium enema for diverticular hemorrhage”

**Mizue Matsuura**, Masahiko Inamori, Atsushi Nakajima, Yasuhiko Komiya, Yumi Inoh, Keigo Kawasima, Mai Naitoh, Yuji Fujita, Akiko Eduka, Noriyoshi Kanazawa, Shiori Uchiyama, Rie Tani, Kennichi Kawana, Setsuya Ohtani, Hajime Nagase.

*World J Gastroenterol.* 2015 May 14;21(18):5555-9. doi: 10.3748/wjg.v21.i18.5555.

### 参考論文

“Feasibility of deep sedation with the combination of propofol and dexmedetomidine hydrochloride for esophageal endoscopic submucosal dissection.”

Nonaka T, Inamori M, Miyashita T, Harada S, Inou Y, Kanoshima K, **Matsuura M**, Higurashi T, Ohkubo H, Iida H, Endo H, Kusakabe A, Maeda S, Gotoh T, Nakajima A.

*Dig Endosc.* 2015 Oct 17.

“Correlation between gastric transit time measured by video capsule endoscopy and gastric emptying determined by the continuous real-time 13C breath test (BreathID system).”

Nonaka T, Inamori M, Endo H, **Matsuura M**, Uchiyama S, Yamada E, Sekino Y, Sakai E, Higurashi T, Ohkubo H, Akimoto K, Iida H, Takahashi H, Koide T, Shibata W, Ida T, Kusakabe A, Gotoh E, Nakajima A, Maeda S.

*Hepatogastroenterology.* 2014 Oct;61(135):2159-62.

“Protective effect of lactoferrin on acute acid reflux-induced esophageal mucosal damage.”

Inamori M, Togawa J, Matsumoto S, Harad K, **Matsuura M**, Iida H, Akimoto K, Endo H, Nonaka T, Takahashi H, Koide T, Shibata W, Ida T, Kusakabe A, Nagase H, Tominaga S, Tanaka K, Gotoh E, Nakajima A, Maeda S.

*Hepatogastroenterology*. 2014 Sep;61(134):1595-600.

“Percutaneous endoscopic gastrostomy in a patient with hemophilia A.”

Koide T, **Iinuma M**, Sakai E, Yamada E, Higurashi T, Ohkubo H, Iida H, Hosono K, Endo H, Nonaka T, Fujita K, Takahashi H, Yoneda M, Goto A, Ida T, Kusakabe A, Nakajima A, Maeda S, Gotoh E, Inamori M.

*Turk J Gastroenterol*. 2014 Dec;25 Suppl 1:241.

“A case of endocrine cell carcinoma of the esophagus successfully treated by chemoradiotherapy.”

Murase T, Higurashi T, Akatsuka S, Nagase H, Kawana K, Otani S, Tani R, Kojima T, Uchiyama S, Eduka A, **Iinuma M**, Fujita Y, Watai K.

*Gan To Kagaku Ryoho*. 2012 Jan;39(1):111-3. Japanese.

“Ulcerative colitis with multidrug-resistant *Pseudomonas aeruginosa* infection successfully treated with bifidobacterium.”

Nagasaki A, Takahashi H, **Iinuma M**, Uchiyama T, Watanabe S, Koide T, Tokoro C, Inamori M, Abe Y, Nakajima A.

*Digestion*. 2010;81(3):204-5. doi: 10.1159/000236042. Epub 2010 Jan 19.